

Regression Lines

Use y on x when:

- x is controlled (not random) variable
- *or* both x and y are random and you are estimating y from given x

- 3 In an agricultural experiment, the relationship between the amount of water supplied, x units, and the yield, y units, was investigated. Six values of x were chosen and for each value of x the corresponding value of y was measured. The results are shown in the table.

x	1	2	3	4	5	6
y	3	6	8	8	11	10

- (i) Give a reason why the regression line of x on y is not suitable in this context. [1]

Binomial & Geometric Distributions – Modelling Assumptions

- For each trial, the probability of a success is constant
- For each trial, the events “success” and “failure” are independent of the corresponding events for every other trial.

These must be contextualised!

They are not the same!

Not a statement about parameter values

- 1 Andy makes repeated attempts to thread a needle. The number of attempts up to and including his first success is denoted by X .

- (i) State two conditions necessary for X to have a geometric distribution. [2]

- (iii) Suggest a reason why one of the conditions you have given in part (i) might not be satisfied in this context. [2]

- 7 At a factory that makes crockery the quality control department has found that 10% of plates have minor faults. These are classed as ‘seconds’. Plates are stored in batches of 12. The number of seconds in a batch is denoted by X .

- (i) State an appropriate distribution with which to model X . Give the value(s) of any parameter(s) and state any assumptions required for the model to be valid. [4]

Poisson Distribution – Modelling Assumptions

- Events occur at constant average rate
- Events occur independently of one another.

These must be contextualised!

“Singly” is a subset of “independently”.

Not a statement about parameter values.

- 5 In a large region of derelict land, bricks are found scattered in the earth.

- (i) State two conditions needed for the number of bricks per cubic metre to be modelled by a Poisson distribution. [2]

Random Sampling

- Every element of the population is equally likely to be chosen
- Each element is chosen independently

People's opinions etc don't have to be independent – it's the selection that matters.

These conditions match those for a binomial distribution.

- 4 A survey is to be carried out to draw conclusions about the proportion p of residents of a town who support the building of a new supermarket. It is proposed to carry out the survey by interviewing a large number of people in the high street of the town, which attracts a large number of tourists.
- (i) Give two different reasons why this proposed method is inappropriate. [2]
- (ii) Suggest a good method of carrying out the survey. [3]
- (iii) State two statistical properties of your survey method that would enable reliable conclusions about p to be drawn. [2]

Central Limit Theorem

- Regardless of the shape of the parent distribution, the mean of a sufficiently large sample is approximately normally distributed

Not a statement about parameter values

Distinguish between “Necessary to use the CLT” and “Possible to use the CLT”.

- 6 The continuous random variable R has the distribution $N(\mu, \sigma^2)$. The results of 100 observations of R are summarised by

$$\Sigma r = 3360.0, \quad \Sigma r^2 = 115\,782.84.$$

- (i) Calculate an unbiased estimate of μ and an unbiased estimate of σ^2 . [4]
- (ii) The mean of 9 observations of R is denoted by \bar{R} . Calculate an estimate of $P(\bar{R} > 32.0)$. [4]
- (iii) Explain whether you need to use the Central Limit Theorem in your answer to part (ii). [2]

Continuity Corrections

- When approximating to a discrete distribution using a continuous

Common: normal approximation to binomial or Poisson

Less common: mean of sample chosen from discrete distribution such as uniform

Here the continuity correction is $\pm 1/2n$, but it's much easier to look at the distribution of the *sum* rather than the *mean*.

- 7 The continuous random variable T is equally likely to take any value from 5.0 to 11.0 inclusive.
- (i) Sketch the graph of the probability density function of T . [2]
- (ii) Write down the value of $E(T)$ and find by integration the value of $\text{Var}(T)$. [5]
- (iii) A random sample of 48 observations of T is obtained. Find the approximate probability that the mean of the sample is greater than 8.3, and explain why the answer is an approximation. [6]