

Kinematics (AS)

Q1	Understand and use the language of kinematics: position; displacement; distance travelled; velocity; speed; acceleration
Q2	Understand, use and interpret graphs in kinematics for motion in a straight line: displacement against time and interpretation of gradient; velocity against time and interpretation of gradient and area under the graph
Q3	Understand, use and derive the formulae for constant acceleration for motion in a straight line

Commentary

Students will already be familiar with vocabulary associated with describing motion. However, it is likely that their understanding of the words, as commonly used in their colloquial language, is not necessarily the same as the definitions of the words when in a technical context. This can present something of a challenge. It is particularly important that students know the difference between a scalar quantity and a vector quantity. It is well worth taking time to ensure that students are clear about the language used to describe motion as this will underpin much of the work ahead.

Use of kinematics graphs is an extremely powerful technique that has been used in various forms at least since the time of Galileo and before the modern practice of using algebraic expressions; they can be used to give information, record information and as a scaffold to guide analysis and interpretation. Students have to understand the conventions used and be sure to use them accurately themselves.

Both the definitions and graphs involve ideas that can be (surprisingly) subtle and it is easy for students to make a false interpretation, especially if the question is phrased ambiguously.

As an example of the need for clarity, suppose Fred is on a trip from Leicester to Manchester and back which is 100 miles each way. How far is Fred from Leicester when he is half way back? Is the answer required 50 miles, 100 miles or 150 miles?

There are some things worth mentioning about the constant acceleration formulae. Of the 5 quantities involved, only one, t , is a scalar. Each of the standard 5 equations connects 4 of the 5 quantities and failure to choose the appropriate equation will lead to inefficient solutions. The processes and results have to be used with care; $v = u + at$ has a starting value, u , so gives v at any time t but the formulae for s give *displacement* over the *elapsed* time t so are not giving position (the standard formulae effectively assume the starting position is 0 when $t = 0$).

Sample MEI resource


'Constant acceleration activity' (found at <http://integralmaths.org/sow-resources.php>) is based on constant acceleration equations. It can be used as game of noughts and crosses. Alternatively students could be asked to find four problems with the same numerical answer.

<p>A particle moving in a straight line is decelerating at a constant rate of 6ms^{-1}.</p> <p>How long will it take to go from a speed of 20ms^{-1} to a speed of 8ms^{-2}?</p>	<p>A book of mass 2kg falls from a shelf 150cm above the floor.</p> <p>Find the speed with which the book hits the floor.</p>	<p>A ball is thrown vertically upwards and takes 3 seconds to reach its highest point.</p> <p>Find the first time the ball is at a height of 39.2m</p>
<p>A train travelling at 35ms^{-1} brakes to a speed of 21ms^{-1} over a distance of 350m.</p> <p>Calculate its deceleration.</p>	<p>A car accelerates from rest at 0.7ms^{-2} for 6s. It then applies its brakes and comes to rest in a distance of 10.5m. Find the total distance travelled and total time taken.</p>	<p>A dog accelerates from rest to 15ms^{-1} in a distance of 30m. Find its acceleration.</p>
<p>A particle is thrown upwards with a velocity of 34.3ms^{-1}.</p> <p>Find how long it takes to reach a height of 49m above the ground.</p>	<p>A stone is dropped from a cliff into the sea below. If the stone hits the water with a speed of 14ms^{-1} find the height of the cliff.</p>	<p>A particle is moving along a straight line with constant accⁿ. It starts from A with velocity of 3ms^{-1} and passes points B and C after 2 and 5 seconds. AC is 40m.</p> <p>Find the acceleration of the particle.</p>

Effective use of technology

'The Moving Man' (which can be found at www.mei.org.uk/integrating-technology) is helpful to get students to see how a simulation can produce various graphs of motion and how they are linked.

The Moving Man



Students should be encouraged to try and simulate various displacement-time and velocity-time graphs. This website can be used to check, discuss their ideas and consider why the graphs produced may be different.

Kinematics(AS)

Time allocation:

Pre-requisites


- Know how to calculate the gradient of a line and the area of regular shapes
- Know the standard units for measuring distance, speed and acceleration
- Solving simple linear and quadratic equations
-

Links with other topics

- Connection to the equation of a straight line and the interpretation of the gradient.
-

Questions and prompts for mathematical thinking

- What is the same and what is different about the motion described by the following?



-

Applications and Modelling

- If you look out of a train window you will see distance markers besides the track every quarter of a mile. Take a train journey and record the time as you go past each marker. How can you use this information to draw distance-time, speed-time and acceleration-time graphs? What can you conclude about the greatest acceleration, deceleration and speed of the train?
-

Common Errors

- Confusing average speed with average velocity
- Assuming negative acceleration always implies deceleration; not appreciating that velocity and acceleration can have different directions
- Not establishing the relevant values, including signs, for the suvat variables.
- Using incorrect formulae; e.g. $v = u + \frac{1}{2}at$, $v^2 = u^2 + 2at$, $s = \left(\frac{v-u}{2}\right)t$.
-