

OCR

Oxford Cambridge and RSA

Wednesday 3 June 2015 – Morning

AS GCE MATHEMATICS (MEI)

4752/01 Concepts for Advanced Mathematics (C2)

QUESTION PAPER

Candidates answer on the Printed Answer Book.


OCR supplied materials:

- Printed Answer Book 4752/01
- MEI Examination Formulae and Tables (MF2)

Other materials required:

- Scientific or graphical calculator

Duration: 1 hour 30 minutes



INSTRUCTIONS TO CANDIDATES

These instructions are the same on the Printed Answer Book and the Question Paper.

- The Question Paper will be found inside the Printed Answer Book.
- Write your name, centre number and candidate number in the spaces provided on the Printed Answer Book. Please write clearly and in capital letters.
- **Write your answer to each question in the space provided in the Printed Answer Book.** Additional paper may be used if necessary but you must clearly show your candidate number, centre number and question number(s).
- Use black ink. HB pencil may be used for graphs and diagrams only.
- Read each question carefully. Make sure you know what you have to do before starting your answer.
- Answer **all** the questions.
- Do **not** write in the bar codes.
- You are permitted to use a scientific or graphical calculator in this paper.
- Final answers should be given to a degree of accuracy appropriate to the context.

INFORMATION FOR CANDIDATES

This information is the same on the Printed Answer Book and the Question Paper.

- The number of marks is given in brackets [] at the end of each question or part question on the Question Paper.
- You are advised that an answer may receive **no marks** unless you show sufficient detail of the working to indicate that a correct method is being used.
- The total number of marks for this paper is **72**.
- The Printed Answer Book consists of **12** pages. The Question Paper consists of **4** pages. Any blank pages are indicated.

INSTRUCTION TO EXAMS OFFICER/INVIGILATOR

- Do not send this Question Paper for marking; it should be retained in the centre or recycled. Please contact OCR Copyright should you wish to re-use this document.

Section A (36 marks)

- 1 (i) Differentiate $12\sqrt[3]{x}$. [2]
- (ii) Integrate $\frac{6}{x^3}$. [3]
- 2 A sequence is defined by $u_1 = 2$ and $u_{k+1} = \frac{10}{u_k^2}$.
Calculate $\sum_{k=1}^4 u_k$. [3]
- 3 An arithmetic progression has tenth term 11.1 and fiftieth term 7.1. Find the first term and the common difference. Find also the sum of the first fifty terms of the progression. [5]
- 4 A sector of a circle has angle 1.5 radians and area 27 cm^2 . Find the perimeter of the sector. [4]
- 5 Use calculus to find the set of values of x for which $x^3 - 6x$ is an increasing function. [5]
- 6 (i) On the same axes, sketch the curves $y = 3^x$ and $y = 3^{2x}$, identifying clearly which is which. [3]
- (ii) Given that $3^{2x} = 729$, find in either order the values of 3^x and x . [2]
- 7 Show that the equation $\sin^2 x = 3\cos x - 2$ can be expressed as a quadratic equation in $\cos x$ and hence solve the equation for values of x between 0 and 2π . [5]
- 8 Fig. 8 shows the graph of $\log_{10} y$ against $\log_{10} x$. It is a straight line passing through the points (2, 8) and (0, 2). [4]




Fig. 8

Find the equation relating $\log_{10} y$ and $\log_{10} x$ and hence find the equation relating y and x . [4]

Section B (36 marks)

9




Fig. 9.1

- (i) Jean is designing a model aeroplane. Fig. 9.1 shows her first sketch of the wing's cross-section. Calculate angle A and the area of the cross-section. [5]
- (ii) Jean then modifies her design for the wing. Fig. 9.2 shows the new cross-section, with 1 unit for each of x and y representing one centimetre.




Fig. 9.2

Here are some of the coordinates that Jean used to draw the new cross-section.

Upper surface		Lower surface	
x	y	x	y
0	0	0	0
4	1.45	4	-0.85
8	1.56	8	-0.76
12	1.27	12	-0.55
16	1.04	16	-0.30
20	0	20	0

Use the trapezium rule with trapezia of width 4cm to calculate an estimate of the area of this cross-section. [6]

- 10 The gradient of a curve is given by $\frac{dy}{dx} = 4x + 3$. The curve passes through the point (2, 9).
- (i) Find the equation of the tangent to the curve at the point (2, 9). [3]
- (ii) Find the equation of the curve and the coordinates of its points of intersection with the x -axis. Find also the coordinates of the minimum point of this curve. [7]
- (iii) Find the equation of the curve after it has been stretched parallel to the x -axis with scale factor $\frac{1}{2}$. Write down the coordinates of the minimum point of the transformed curve. [3]
- 11 Jill has 3 daughters and no sons. They are generation 1 of Jill's descendants.

Each of her daughters has 3 daughters and no sons. Jill's 9 granddaughters are generation 2 of her descendants. Each of her granddaughters has 3 daughters and no sons; they are descendant generation 3.

Jill decides to investigate what would happen if this pattern continues, with each descendant having 3 daughters and no sons.

- (i) How many of Jill's descendants would there be in generation 8? [2]
- (ii) How many of Jill's descendants would there be altogether in the first 15 generations? [3]
- (iii) After n generations, Jill would have over a million descendants altogether. Show that n satisfies the inequality

$$n > \frac{\log_{10} 2000003}{\log_{10} 3} - 1.$$

Hence find the least possible value of n . [4]

- (iv) How many **fewer** descendants would Jill have altogether in 15 generations if instead of having 3 daughters, she and each subsequent descendant has 2 daughters? [3]

END OF QUESTION PAPER

Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact the Copyright Team, First Floor, 9 Hills Road, Cambridge CB2 1GE.

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

OCR

Oxford Cambridge and RSA

Wednesday 3 June 2015 – Morning

AS GCE MATHEMATICS (MEI)

4752/01 Concepts for Advanced Mathematics (C2)

PRINTED ANSWER BOOK

Candidates answer on this Printed Answer Book.

OCR supplied materials:

- Question Paper 4752/01 (inserted)
- MEI Examination Formulae and Tables (MF2)

Other materials required:

- Scientific or graphical calculator

Duration: 1 hour 30 minutes



Candidate forename		Candidate surname	
--------------------	--	-------------------	--

Centre number						Candidate number				
---------------	--	--	--	--	--	------------------	--	--	--	--

INSTRUCTIONS TO CANDIDATES

These instructions are the same on the Printed Answer Book and the Question Paper.

- The Question Paper will be found inside the Printed Answer Book.
- Write your name, centre number and candidate number in the spaces provided on the Printed Answer Book. Please write clearly and in capital letters.
- **Write your answer to each question in the space provided in the Printed Answer Book.** Additional paper may be used if necessary but you must clearly show your candidate number, centre number and question number(s).
- Use black ink. HB pencil may be used for graphs and diagrams only.
- Read each question carefully. Make sure you know what you have to do before starting your answer.
- Answer **all** the questions.
- Do **not** write in the bar codes.
- You are permitted to use a scientific or graphical calculator in this paper.
- Final answers should be given to a degree of accuracy appropriate to the context.

INFORMATION FOR CANDIDATES

This information is the same on the Printed Answer Book and the Question Paper.

- The number of marks is given in brackets [] at the end of each question or part question on the Question Paper.
- You are advised that an answer may receive **no marks** unless you show sufficient detail of the working to indicate that a correct method is being used.
- The total number of marks for this paper is **72**.
- The Printed Answer Book consists of **12** pages. The Question Paper consists of **4** pages. Any blank pages are indicated.

Section A (36 marks)

1 (i)	
1 (ii)	
2	
3	

4

5

6(i)**6(ii)**

7

8

Section B (36 marks)

9(i)	

9(ii)

Upper surface		Lower surface	
x	y	x	y
0	0	0	0
4	1.45	4	-0.85
8	1.56	8	-0.76
12	1.27	12	-0.55
16	1.04	16	-0.30
20	0	20	0

10 (i)	
10 (ii)	
(answer space continued on next page)	

10 (ii)	(continued)
10 (iii)	

11 (i)	
11 (ii)	

11 (iii)	
11 (iv)	

PLEASE DO NOT WRITE ON THIS PAGE



Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact the Copyright Team, First Floor, 9 Hills Road, Cambridge CB2 1GE.

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

Annotations and abbreviations

Annotation in scoris	Meaning
✓ and ✖	
BOD	Benefit of doubt
FT	Follow through
ISW	Ignore subsequent working
M0, M1	Method mark awarded 0, 1
A0, A1	Accuracy mark awarded 0, 1
B0, B1	Independent mark awarded 0, 1
SC	Special case
^	Omission sign
MR	Misread
Highlighting	
Other abbreviations in mark scheme	Meaning
E1	Mark for explaining
U1	Mark for correct units
G1	Mark for a correct feature on a graph
M1 dep*	Method mark dependent on a previous mark, indicated by *
cao	Correct answer only
oe	Or equivalent
rot	Rounded or truncated
soi	Seen or implied
www	Without wrong working

Subject-specific Marking Instructions for GCE Mathematics (MEI) Pure strand

- a Annotations should be used whenever appropriate during your marking.

The A, M and B annotations must be used on your standardisation scripts for responses that are not awarded either 0 or full marks. It is vital that you annotate standardisation scripts fully to show how the marks have been awarded.

For subsequent marking you must make it clear how you have arrived at the mark you have awarded.

- b An element of professional judgement is required in the marking of any written paper. Remember that the mark scheme is designed to assist in marking incorrect solutions. Correct *solutions* leading to correct answers are awarded full marks but work must not be judged on the answer alone, and answers that are given in the question, especially, must be validly obtained; key steps in the working must always be looked at and anything unfamiliar must be investigated thoroughly.

Correct but unfamiliar or unexpected methods are often signalled by a correct result following an *apparently* incorrect method. Such work must be carefully assessed. When a candidate adopts a method which does not correspond to the mark scheme, award marks according to the spirit of the basic scheme; if you are in any doubt whatsoever (especially if several marks or candidates are involved) you should contact your Team Leader.

- c The following types of marks are available.

M

A suitable method has been selected and *applied* in a manner which shows that the method is essentially understood. Method marks are not usually lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, eg by substituting the relevant quantities into the formula. In some cases the nature of the errors allowed for the award of an M mark may be specified.

A

Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated Method mark is earned (or implied). Therefore M0 A1 cannot ever be awarded.

B

Mark for a correct result or statement independent of Method marks.

E

A given result is to be established or a result has to be explained. This usually requires more working or explanation than the establishment of an unknown result.

Unless otherwise indicated, marks once gained cannot subsequently be lost, eg wrong working following a correct form of answer is ignored. Sometimes this is reinforced in the mark scheme by the abbreviation isw. However, this would not apply to a case where a candidate passes through the correct answer as part of a wrong argument.

- d When a part of a question has two or more 'method' steps, the M marks are in principle independent unless the scheme specifically says otherwise; and similarly where there are several B marks allocated. (The notation 'dep **' is used to indicate that a particular mark is dependent on an earlier, asterisked, mark in the scheme.) Of course, in practice it may happen that when a candidate has once gone wrong in a part of a question, the work from there on is worthless so that no more marks can sensibly be given. On the other hand, when two or more steps are successfully run together by the candidate, the earlier marks are implied and full credit must be given.
- e The abbreviation ft implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A and B marks are given for correct work only — differences in notation are of course permitted. A (accuracy) marks are not given for answers obtained from incorrect working. When A or B marks are awarded for work at an intermediate stage of a solution, there may be various alternatives that are equally acceptable. In such cases, exactly what is acceptable will be detailed in the mark scheme rationale. If this is not the case please consult your Team Leader.

Sometimes the answer to one part of a question is used in a later part of the same question. In this case, A marks will often be 'follow through'. In such cases you must ensure that you refer back to the answer of the previous part question even if this is not shown within the image zone. You may find it easier to mark follow through questions candidate-by-candidate rather than question-by-question.

- f Wrong or missing units in an answer should not lead to the loss of a mark unless the scheme specifically indicates otherwise. Candidates are expected to give numerical answers to an appropriate degree of accuracy, with 3 significant figures often being the norm. Small variations in the degree of accuracy to which an answer is given (e.g. 2 or 4 significant figures where 3 is expected) should not normally be penalised, while answers which are grossly over- or under-specified should normally result in the loss of a mark. The situation regarding any particular cases where the accuracy of the answer may be a marking issue should be detailed in the mark scheme rationale. If in doubt, contact your Team Leader.
- g Rules for replaced work

If a candidate attempts a question more than once, and indicates which attempt he/she wishes to be marked, then examiners should do as the candidate requests.

If there are two or more attempts at a question which have not been crossed out, examiners should mark what appears to be the last (complete) attempt and ignore the others.

NB Follow these maths-specific instructions rather than those in the assessor handbook.

- h For a *genuine* misreading (of numbers or symbols) which is such that the object and the difficulty of the question remain unaltered, mark according to the scheme but following through from the candidate's data. A penalty is then applied; 1 mark is generally appropriate, though this may differ for some units. This is achieved by withholding one A mark in the question.

Note that a miscopy of the candidate's own working is not a misread but an accuracy error.

Question		Answer	Marks	Guidance
1	(i)	$kx^{\frac{1}{3}-1}$ oe $4x^{\frac{-2}{3}}$ isw cao	M1 A1 [2]	k is any non-zero constant ignore $+c$ allow any equivalent exact simplified form
1	(ii)	kx^{-3+1} oe $-3x^{-2}$ isw $+c$	M1 A1 A1 [3]	k is any non-zero constant allow any equivalent exact simplified form
2		$u_2 = \frac{10}{2^2}, u_3 = \frac{10}{\text{their } 2.5^2}, u_4 = \frac{10}{\text{their } 1.6^2}$ isw $2 + u_2 + u_3 + u_4$ soi 10.00625 or $\frac{1601}{160}$ or $10\frac{1}{80}$ cao isw	M1* M1dep* A1 [3]	NB 2.5, 1.6, 3.90625 or $\frac{10}{4}, \frac{8}{5}, \frac{125}{32}$ may be implied by eg sight of 3.9 and answer of 10.0 NB 2.5, 1.1, 0.625 scores M0M0 B3 if unsupported

3		$a + (10 - 1)d = 11.1$ and $a + (50 - 1)d = 7.1$ $d = -0.1$ $a = 12$ $\frac{1}{2} \times 50(\text{their } a + 7.1)$ with $a > 11.1$ 477.5 or $477\frac{1}{2}$ or $\frac{955}{2}$ cao	M1 A1 A1 M1 A1 [5]	may be implied by $40d = \pm 4$ or embedded in attempt to solve if unsupported, B2 for one of these and B3 for both or $\frac{50}{2}(2a + (50 - 1)d)$ with $a > 11.1$ and $d < 0$	condone one slip in coefficient of d if M0, B2 for any form of correct answer www
4		$27 = \frac{1}{2} r^2 \times 1.5$ oe $r = 6$ soi their $r \times 1.5$ 21 [cm] cao	M1 A1 M1 A1 [4]	or $27 = \frac{85.943669\dots}{360} \times \pi r^2$ may be embedded in formula for arc length or their $\frac{85.943639}{360} \times 2\pi \times \text{their } r$ allow full marks for recovery from working with rounded value of θ in degree form	angle in degrees rounded to 2 sf or more may be implied by later work eg 9 or 21 if r is incorrect, we must see their $r \times 1.5 [+ 2r]$ for M1 if r is correct, M1 may be implied by 9 or 21 B4 for 21 unsupported www

5		$3x^2 - 6$ seen <i>their</i> $y' = 0$ or $y' > 0$ or $y' \geq 0$ $\sqrt{2}$ and $-\sqrt{2}$ identified $x < -\sqrt{2}$ or $x \leq -\sqrt{2}$ isw $x > \sqrt{2}$ or $x \geq \sqrt{2}$	B1 M1 A1 A1 A1 [5]	must be quadratic with at least one of only two terms correct may be implied by use with inequalities or by $\pm 1.41[4213562]$ to 3 sf or more if A1A0A0 , allow SC1 for fully correct answer in decimal form to 3 sf or more or A2 for $ x > \sqrt{2}$ or $ x \geq \sqrt{2}$	$ x = \sqrt{2}$ implies A1 NB just $-\sqrt{2} > x > \sqrt{2}$ or $\sqrt{2} < x < -\sqrt{2}$ or $x > \pm\sqrt{2}$ implies the first A1 then A0A0
6	(i)	both curves with positive gradients in 1 st and 2 nd quadrants; ignore labels for this mark both through (0, 1) $y = 3^{2x}$ above $y = 3^x$ in first quadrant and below it in second	M1 A1 A1 [3]	do not award if clearly not exponential shape; condone touching negative x -axis but not crossing it must be clearly labelled, A0 if wrongly attributed or if coincide for negative x from (0, 1)	consider each curve independently; ignore scales and points apart from (0, 1) allow if indicated in table of values or commentary if not marked on graph if M0 allow SC1 for one graph fully correct
6	(ii)	$x = 3$ $3^x = 27$	B1 B1 [2]	B0 if wrongly attributed B0 if wrongly attributed	allow $3^3 = 27$ with $x = 3$ stated

7		$1 - \cos^2 x = 3\cos x - 2$ oe $\cos^2 x + 3\cos x - 3 [= 0]$ $\cos x = \text{their } \frac{-3 + \sqrt{21}}{2}$ or $\cos x = \text{their } 0.79 \text{ to } 0.7913$ soi $[x =] 0.6578 \text{ to } 0.66$ isw cao $[x =] 5.625 \text{ to } 5.63$ isw cao	M1* M1*dep M1 A1 A1 [5]	or $-\cos^2 x - 3\cos x + 3 = 0$ dependent on award of previous method mark, must be correct for their quadratic A0 for eg 0.66π if 0.66 not seen separately if A1A1 extra values in range incur a penalty of 1; ignore extra values outside range if A0A0 allow SC1 for 37.69 to 37.7° and 322 to 322.31° or for $(0.209 \text{ to } 0.21)\pi$ and $(1.79 \text{ to } 1.791)\pi$	condone one sign error <i>or</i> constant term of -1 (in LH version) or $+1$ (in RH version) ignore other values (eg $-3.79\dots$); condone recovery from $x = 0.791287847\dots$ but M0 if no recovery NB $x = 0.65788395\dots$ NB $x = 5.625301357\dots$ no SC mark available if extra values in range
8		$m = 3$ seen $\log y = m\log x + 2$ or $\log y = m\log x + \log 100$ $\log y = \log x^3 + 2$ or $\log y = \log x^3 + \log 100$ or better $y = 100x^3$ or $y = 10^{3\log x + 2}$ or $y = 10^{\log x^3 + 2}$ www isw	B1 M1 M1 A1 [4]	or $\log y - 8 = m(\log x - 2)$ or $10^{\log y} = 10^{3\log x + 2}$ or $10^{3\log x + \log 100}$ or better $y = 10^{3\log x + \log 100}$ or $y = 10^{\log x^3 + \log 100}$	condone lack of base; “ $c = 2$ ” is insufficient condone lack of base, but not bases other than 10 unless fully recovered

9	(i)	<p>$[\cos A =] \frac{20^2 + 13^2 - 8^2}{2 \times 13 \times 20}$</p> <p>$[\cos A =] \frac{505}{520}$ oe soi</p> <p>$A = 13.79$ to 13.8° or 14°</p> <p>$[\text{Area} =] \frac{1}{2} \times 20 \times 13 \times \sin \text{their } A$</p> <p>$30.99$ to 31.01 isw</p> <p>or $\frac{5\sqrt{615}}{4}$ oe isw</p>	<p>M1*</p> <p>A1</p> <p>A1</p> <p>M1dep*</p> <p>A1</p> <p>[5]</p>	<p>or $8^2 = 20^2 + 13^2 - 2 \times 13 \times 20 \times \cos A$</p> <p>or 0.971 to 0.9712</p> <p>or 0.24077 to 0.241 or 0.24 (radians); allow B3 if given to 3sf or more unsupported</p> <p>or M1 for eg $\frac{1}{2} \times 20 \times 8 \times \sin 22.8$, as long as angle calculated correctly from their A (other angles are $22.79824\dots^\circ$ and $143.40645\dots^\circ$ or $36.59355\dots^\circ$)</p> <p>allow B2 for unsupported answer within range</p>	<p>or 15.32 (grad)</p> <p>or</p> $\sqrt{\frac{41}{2} \left(\frac{41}{2} - 8\right) \left(\frac{41}{2} - 13\right) \left(\frac{41}{2} - 20\right)}$ <p>NB $13\sin A = 3.099899192$ if $\frac{1}{2} \times b \times h$ used</p>
---	-----	---	---	---	--

9	(ii)	<p>$h = 4$ soi</p> <p>$\frac{\text{their } 4}{2} \times (0 + 0 + 2(1.45+1.56+1.27+1.04))$</p> <p>or</p> <p>$\frac{\text{their } 4}{2} \times (0 + 0 + 2(\pm 0.85 \pm 0.76 \pm 0.55 \pm 0.30))$</p> <p>either 21.28 or ± 9.84</p> <p>their 21.28 + their 9.84</p> <p>31.12</p>	<p>B1</p> <p>M1*</p> <p>B1</p> <p>A1</p> <p>M1dep*</p> <p>A1</p>	<p>shape of formula correct with 2, 3 or 4 y-values in inner bracket with their h; allow recovery from bracket errors</p> <p>M0 if any non-zero x-values used or if y-values used twice</p> <p>all y-values correctly placed with their h, condone omission of zeros and/or omission of outer brackets</p> <p>ignore subsequent rounding, but A0 if answer spoiled by eg multiplication by 20</p>	<p>eg $\frac{\text{their } 4}{2} \times \{1.45 + 1.04 + 2(1.56 + 1.27)\}$; signs must be consistent in 2nd alternative</p> <p>or B1 + B3* if area of 2 triangles and 3 trapezia calculated to give correct answer www The final M1dep* A1 may then be earned.</p> <p>NB $2.9 + 6.02 + 5.66 + 4.62 + 2.08$ or $\pm 1.7 \pm 3.22 \pm 2.62 \pm 1.7 \pm 0.60$ with consistent signs throughout</p>
---	------	---	--	--	--

9	(ii)	<p><i>alternatively</i></p> <p>$h = 4$ soi</p> <p>attempt to find all y-values</p> <p>2.3, 2.32, 1.82, 1.34</p> <p>$\frac{\text{their } 4}{2} \times (0 + 0 + 2(2.3+2.32+1.82+1.34))$</p> <p>31.12</p>	<p>B1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>B1FT</p> <p>A1</p> <p>[6]</p>	<p>$y_{\text{upper}} - y_{\text{lower}}$</p> <p>all y-values correct</p> <p>shape of formula correct with 2, 3 or 4 of their y-values in inner bracket with their h; allow recovery from bracket errors</p> <p>M0 if any non-zero x-values used or if y-values used twice</p> <p>all their y-values correctly placed, condone omission of zeros and/or omission of outer brackets</p> <p>ignore subsequent rounding, but A0 if answer spoiled by eg multiplication by 20</p>	<p>M0 if values are added to obtain 0.60, 0.80 etc</p> <p>eg $\frac{1}{2} \times 4 \times \{2.3 + 1.34 + 2(2.32+1.82)\}$</p> <p>or B1M1A1 + B3 if area of 2 triangles and 3 trapezia calculated to give correct answer www NB $4.6 + 9.24 + 8.28 + 6.32 + 2.68$</p>
---	------	---	---	---	---

10	(i)	$\left[\frac{dy}{dx}\right] 4 \times 2 + 3 \text{ or } 11 \text{ isw}$ $9 = \text{their } (4 \times 2 + 3) \times 2 + c$ $y = 11x - 13 \text{ or } y = 11x + c \text{ and } c = -13$ <p>stated isw</p>	<p>M1*</p> <p>M1dep*</p> <p>A1</p> <p>[3]</p>	<p>or $y - 9 = \text{their } (4 \times 2 + 3) \times (x - 2)$</p> <p>or $y - 9 = 11(x - 2)$ isw</p>	
10	(ii)	$\frac{4x^2}{2} + 3x$ $[y =] 2x^2 + 3x + c$ $9 = 2 \times 2^2 + 3 \times 2 + c$ $y = 2x^2 + 3x - 5 \text{ cao}$ <p>$(1, 0)$ and $(-2.5, 0)$ oe cao</p> $x = -\frac{3}{4}$ $y = -\frac{49}{8}$	<p>M1*</p> <p>A1</p> <p>M1dep*</p> <p>A1</p> <p>B1</p> <p>B1</p> <p>B1</p> <p>[7]</p>	<p>must see “2” and “+ c”; may be earned later eg after attempt to find c</p> <p>must include constant, which may be implied by answer</p> <p>allow first 4 marks for $y = 2x^2 + 3x + c$ and $c = -5$ stated</p> <p>or for $x = 1, y = 0$ and $x = -2.5, y = 0$</p> <p>-6.125 or $-6\frac{1}{8}$</p>	<p>B0 for just stating $x = 1$ and $x = -2.5$</p>

10	(iii)	substitution to obtain $[y =] f(2x)$ in polynomial form $y = (2x - 1)(4x + 5)$ or $y = 8x^2 + 6x - 5$ or $y = 2\left(2x + \frac{3}{4}\right)^2 - \frac{49}{8}$ $\left(-\frac{3}{8}, -\frac{49}{8}\right)$ oe	M1 A1FT B1 [3]	$f(x)$ must be the quadratic in x with linear and constant term obtained in part (ii), may be in factorised form must be simplified to one of these forms, FT their quadratic in x with linear and constant term obtained in part (ii) or FT their (both non-zero) co-ordinates for minimum point or their quadratic in x with linear and constant term obtained in part (ii)	or their $x = 1 \rightarrow$ their 0.5 and their $x = -2.5 \rightarrow$ their $x = -1.25$ hence $y = (2x - 1)(4x + 5)$ FT their x -intercepts from their quadratic in x with linear and constant term obtained in part (ii)
11	(i)	3×3^7 oe 6561	M1 A1 [2]	condone 1×3^7 or B2 if unsupported	do not award if only seen in sum of terms of GP if 0, SC1 for 2187 unsupported
11	(ii)	valid attempt to sum a GP with $r = 3$ and $n = 15$ $\frac{3(3^{15} - 1)}{3 - 1}$ oe 21 523 359	M1 M1 A1 [3]	eg $3 + 3^2 + \dots + 3^{15}$ or B2 if M1M0 or B3 if unsupported	must see at least first two terms and last term NB 7 174 453 implies M1 from $1 + 3 + \dots + 3^{14}$

11	(iii)	$\frac{3(3^n - 1)}{3 - 1} > 1000000$ <p>eg $3^{n+1} > 2000003$ or $3^n > \frac{2000000}{3} + 1$</p> <p>www</p> <p>correctly taking logs of both sides</p> <p>eg $(n + 1) \log 3 > \log 2000003$ or $n \log 3 > \log 2000003 - \log 3$</p> <p>eg $n + 1 > \frac{\log 2000003}{\log 3}$ and completion to</p> $n > \frac{\log 2000003}{\log 3} - 1$ <p>$n = 13$ seen</p>	<p>M1*</p> <p>M1dep*</p> <p>A1</p> <p>B1</p> <p>[4]</p>	<p>eg $\log 3^{n+1} > \log 2000003$ www or $\log 3^n + \log 3 > \log 2000003$ www; may be implied by next stage of working</p> <p>without any wrong working</p> <p>B0 for $n \geq 13$ or $n > 13$</p>	<p>M0 for working backwards</p> <p>M0 if = or < used</p> <p>at least one previous progressive interim step needed with no wrong working; M0dep* for $\log(3^n - 1) > \dots$</p> <p>do not allow recovery from bracket errors at any stage</p>
11	(iv)	<p>valid attempt to sum a GP with $r = 2$ and $n = 15$</p> <p>their 21 523 359 – their 65 534 21 457 825 isw</p>	<p>M1*</p> <p>M1dep*</p> <p>A1</p> <p>[3]</p>	<p>if correct eg $2 + 2^2 + \dots + 2^{15} = 65\,534$</p> <p>with their $65\,534 <$ their $21\,523\,359$</p> <p>allow B3 for 21 457 825 unsupported</p>	<p>NB 32767 implies M1 from $1 + 2 + \dots + 2^{14}$</p>

GCE Mathematics (MEI)			Max Mark	a	b	c	d	e	u
4751	01 C1 – MEI Introduction to advanced mathematics (AS)	Raw	72	63	58	53	48	43	0
		UMS	100	80	70	60	50	40	0
4752	01 C2 – MEI Concepts for advanced mathematics (AS)	Raw	72	56	50	44	39	34	0
		UMS	100	80	70	60	50	40	0
4753	01 (C3) MEI Methods for Advanced Mathematics with Coursework: Written Paper	Raw	72	56	51	46	41	36	0
4753	02 (C3) MEI Methods for Advanced Mathematics with Coursework: Coursework	Raw	18	15	13	11	9	8	0
4753	82 (C3) MEI Methods for Advanced Mathematics with Coursework: Carried Forward Coursework Mark	Raw	18	15	13	11	9	8	0
		UMS	100	80	70	60	50	40	0
4754	01 C4 – MEI Applications of advanced mathematics (A2)	Raw	90	74	67	60	54	48	0
		UMS	100	80	70	60	50	40	0
4755	01 FP1 – MEI Further concepts for advanced mathematics (AS)	Raw	72	62	57	53	49	45	0
		UMS	100	80	70	60	50	40	0
4756	01 FP2 – MEI Further methods for advanced mathematics (A2)	Raw	72	65	58	52	46	40	0
		UMS	100	80	70	60	50	40	0
4757	01 FP3 – MEI Further applications of advanced mathematics (A2)	Raw	72	59	52	46	40	34	0
		UMS	100	80	70	60	50	40	0
4758	01 (DE) MEI Differential Equations with Coursework: Written Paper	Raw	72	63	57	51	45	38	0
4758	02 (DE) MEI Differential Equations with Coursework: Coursework	Raw	18	15	13	11	9	8	0
4758	82 (DE) MEI Differential Equations with Coursework: Carried Forward Coursework Mark	Raw	18	15	13	11	9	8	0
		UMS	100	80	70	60	50	40	0
4761	01 M1 – MEI Mechanics 1 (AS)	Raw	72	62	54	46	39	32	0
		UMS	100	80	70	60	50	40	0
4762	01 M2 – MEI Mechanics 2 (A2)	Raw	72	54	47	40	33	27	0
		UMS	100	80	70	60	50	40	0
4763	01 M3 – MEI Mechanics 3 (A2)	Raw	72	64	56	48	41	34	0
		UMS	100	80	70	60	50	40	0
4764	01 M4 – MEI Mechanics 4 (A2)	Raw	72	53	45	38	31	24	0
		UMS	100	80	70	60	50	40	0
4766	01 S1 – MEI Statistics 1 (AS)	Raw	72	61	54	47	41	35	0
		UMS	100	80	70	60	50	40	0
4767	01 S2 – MEI Statistics 2 (A2)	Raw	72	65	60	55	50	46	0
		UMS	100	80	70	60	50	40	0
4768	01 S3 – MEI Statistics 3 (A2)	Raw	72	64	58	52	47	42	0
		UMS	100	80	70	60	50	40	0
4769	01 S4 – MEI Statistics 4 (A2)	Raw	72	56	49	42	35	28	0
		UMS	100	80	70	60	50	40	0
4771	01 D1 – MEI Decision mathematics 1 (AS)	Raw	72	56	51	46	41	37	0
		UMS	100	80	70	60	50	40	0
4772	01 D2 – MEI Decision mathematics 2 (A2)	Raw	72	54	49	44	39	34	0
		UMS	100	80	70	60	50	40	0
4773	01 DC – MEI Decision mathematics computation (A2)	Raw	72	46	40	34	29	24	0
		UMS	100	80	70	60	50	40	0
4776	01 (NM) MEI Numerical Methods with Coursework: Written Paper	Raw	72	56	50	45	40	34	0
4776	02 (NM) MEI Numerical Methods with Coursework: Coursework	Raw	18	14	12	10	8	7	0
4776	82 (NM) MEI Numerical Methods with Coursework: Carried Forward Coursework Mark	Raw	18	14	12	10	8	7	0
		UMS	100	80	70	60	50	40	0
4777	01 NC – MEI Numerical computation (A2)	Raw	72	55	47	39	32	25	0
		UMS	100	80	70	60	50	40	0
4798	01 FPT - Further pure mathematics with technology (A2)	Raw	72	57	49	41	33	26	0
		UMS	100	80	70	60	50	40	0

GCE Statistics (MEI)										
			Max Mark	a	b	c	d	e	u	
G241	01	Statistics 1 MEI (Z1)	Raw	72	61	54	47	41	35	0
			UMS	100	80	70	60	50	40	0
G242	01	Statistics 2 MEI (Z2)	Raw	72	55	48	41	34	27	0
			UMS	100	80	70	60	50	40	0
G243	01	Statistics 3 MEI (Z3)	Raw	72	56	48	41	34	27	0
			UMS	100	80	70	60	50	40	0

GCE Quantitative Methods (MEI)										
			Max Mark	a	b	c	d	e	u	
G244	01	Introduction to Quantitative Methods MEI	Raw	72	58	50	43	36	28	0
G244	02	Introduction to Quantitative Methods MEI	Raw	18	14	12	10	8	7	0
			UMS	100	80	70	60	50	40	0
G245	01	Statistics 1 MEI	Raw	72	61	54	47	41	35	0
			UMS	100	80	70	60	50	40	0
G246	01	Decision 1 MEI	Raw	72	56	51	46	41	37	0
			UMS	100	80	70	60	50	40	0